

## **Yam Ointment**

**For external veterinary use**

### **Dosage Form**

**Ointment for external use in veterinary medicine.**

### **Pharmacotherapeutic Group**

**Fungicidal and bactericidal medicinal product.**

### **Target Species**

**Animals.**

---

### **Composition**

**1 g of the product contains active substances:**

- **Zinc oxide – 100 mg**
  - **Sulfur – 100 mg**
  - **Salicylic acid – 20 mg**
- 

### **Description**

**Gray or gray to brown ointment with a specific odor.**

**Layer separation of components may occur during prolonged storage.**

---

### **Pharmacological Properties**

**Combined preparation for external use.**

**It has pronounced fungicidal, bactericidal and ectoparasitic properties due to the synergistic action of its components.**

**The active ingredients ensure penetration into the upper layers of the skin and destroy pathogens deep within the lesion site.**

**The product is slightly toxic to warm-blooded animals and does not exhibit irritant or sensitizing effects.**

---

## **Indications**

**Used for the treatment of various skin diseases in animals, including:**

- **Eczema**
- **Dermatitis**
- **Trichophytosis**
- **Other fungal and bacterial skin diseases**

**Also effective in:**

- **Hoof lesions**
- **Necrobacteriosis**
- **Sheep foot rot**
- **Sarcoptic mange (scabies)**

---

## **Dosage and Administration**

**Before use, the ointment should be thoroughly mixed, as prolonged storage may cause separation of components.**

**Apply a thin layer to the affected area and 2–4 cm around it.**

**Treat the lesion 1–2 times daily.**

**If scabs do not detach within 4–5 days, continue treatment for 7–10 days.**

**Hair regrowth and clearing of the affected area are usually observed.**

**After 10 days, microscopic examination of the affected skin is recommended.**

**If the pathogen is detected, the treatment course should be repeated.**

---

## **Adverse Reactions**

**No adverse reactions have been observed when used as directed.**

**Hypersensitivity to product components may occur.**

---

**Contraindications**

None established.

---

**Special Instructions**

Observe personal hygiene and safety regulations when handling veterinary medicinal products.

---

**Storage Conditions**

Store in a dry place protected from light at a temperature not exceeding 25°C.

---

**Presentation**

20 g and 0.5 kg ointment in polyethylene containers.

---

**Shelf Life**

2 years.

---

**Manufacturer**

LLC "Biotecs"  
Tbilisi, Georgia  
8 Yumashevi Street